


## Information for patients



### Posterior Vitreous Detachment



### An explanation of causes and treatment


A clear jelly, (the vitreous or 'vitreous jelly'), fills the middle of the eye. The vitreous lies against the retina in the normal eye. It is transparent like glass, so light passes through it to reach the retina: the retina is the film that lines the back of the eye.

#### The Vitreous



## How the vitreous changes

As you get older the vitreous may shrink away from the retina. This may happen earlier if you are short sighted or have injured your eye. This shrinking process may happen rather suddenly, that is over a few days. This process is called a posterior vitreous detachment. The eye still sees well with a shrunken vitreous:




The shrinkage is essentially like a jelly liquefying, and no harm comes to the eye. The vitreous may shrink in different ways.

First, it may shrink away from the retina, and leave the retina unaffected. You may not notice if this happens.

Secondly, it may tug the retina gently. This may cause tiny flashes of light. These usually subside over a couple of weeks.

Thirdly, you may develop floaters. These are vitreous opacities that cast a shadow on the retina.


### Floaters



You may see these as a spider's web or veil over the eye. Floaters disappear a little, and become less noticeable, over the next few weeks or months.

Floaters are naturally much more noticeable if you only have one good eye (and this process is happening in the good eye).

Less commonly, the vitreous may pull the retina and make a small retinal tear, or even less commonly, a detached retina.



## **The Eye Examination**

In the eye clinic or eye emergency department your eye will be examined. So the doctor can see the edge of the retina to look for a retinal tear, your pupil will be dilated with drops. The drops take about 20 minutes to work, and your reading sight will be blurred for about two hours. (very occasionally the sight is slightly blurred for a day or two.)

The examination may be carried out with several types of instrument, or even a contact lens, and occasionally the doctor has to press on the edge of the eye. This may be a little painful.

## **The treatment - none usually**

There is no treatment that will put the vitreous back in position. The floaters and veil that may have drifted across your sight subside by themselves. You may notice a large floater for a long time, which is a nuisance; the doctor can not remove this. As mentioned, most people became accustomed to the floater or floaters, and with a little effort ignore them.

## **A Retinal Tear**

Tears are uncommon, but if one develops, laser treatment around it, or freezing therapy, may be necessary to prevent the tear becoming larger.

## **Precautions**

If the doctor checks your eye and all is well, the floaters and flashers subside.

However, you usually need a further examination if:

- You suddenly develop a lot more flashes or floaters (this could be a small tear)
- A shutter or curtain of blurred vision drifts across your eye, sometimes from below. To check for this, cover one eye at a time for a few seconds every day. If the eye you have not covered sees well in all directions, all is well.
- For advice contact: Victoria Eye Unit, the County Hospital, Hereford Tel: 01432 355 444 ext 1766

Issue Date: March 2008

Review Date: November 2014